

*Leave answers in simplest radical or fractional form. NO DECIMAL ANSWERS!!

I. Sum or Difference Identities.

A. Find the exact value of each trig value by rewriting the angle as the sum or difference of 2 special angles.

1. $\cos \frac{\pi}{12}$

2. $\sin 105^\circ$

3. $\tan 15^\circ$

B. Find the exact value of each expression by simplifying to a special angle.

4. $\cos 40^\circ \cos 10^\circ + \sin 40^\circ \sin 10^\circ$

5. $\sin \frac{\pi}{12} \cos \frac{7}{12} \pi - \cos \frac{\pi}{12} \sin \frac{7}{12} \pi$

6. $\frac{\tan 40^\circ + \tan 20^\circ}{1 - \tan 40^\circ \tan 20^\circ}$

II. Double and Half Angle Identities.

A. Find the exact value of each trig ratio by rewriting the angle as half of a special angle.

7. $\sin 195^\circ$

8. $\cos 165^\circ$

9. $\tan \frac{9}{8} \pi$

B. Given: $\sin A = -\frac{3}{5}$, $\pi < A < \frac{3}{2}\pi$ and $\cos B = -\frac{12}{13}$, $\frac{\pi}{2} < B < \pi$

Set up a triangle in the correct quadrant for angle A and angle B. Use the triangle to find each value.

Angle A

Angle B

10. $\sin(A - B)$

11. $\tan(A + B)$

12. $\cos 2B$

13. $\cos \frac{A}{2}$

Verify each identity. Use ONLY the Fundamental Identities. Show ALL Steps.

14. $3\sin^2\theta + 4\cos^2\theta = 3 + \cos^2\theta$

15. $\frac{\sec x}{\csc x} + \frac{\sin x}{\cos x} = 2 \tan x$